

SAMHSA
Opioid Overdose
TOOLKIT:
Information for Prescribers



TABLE OF CONTENTS

INFORMATION FOR PRESCRIBERS

OPIOID OVERDOSE	3
TREATING OPIOID OVERDOSE	7
LEGAL AND LIABILITY CONSIDERATIONS	9
CLAIMS CODING AND BILLING	9
RESOURCES FOR PRESCRIBERS	9
ACKNOWLEDGMENTS, ETC.	11
■ Acknowledgments	
■ Disclaimer	
■ Public Domain Notice	
■ Electronic Access and Copies of Publication	
■ Recommended Citation	
■ Originating Office	

Also see the other components of this Toolkit:

- ★ Facts for Community Members
- ★ Five Essential Steps for First Responders
- ★ Safety Advice for Patients & Family Members
- ★ Recovering from Opioid Overdose:
Resources for Overdose Survivors & Family Members

Opioid overdose is a major public health problem, accounting for almost 17,000 deaths a year in the United States [1]. Overdose involves both males and females of all ages, ethnicities, and demographic and economic characteristics, and involves both illicit opioids such as heroin and, increasingly, prescription opioid analgesics such as oxycodone, hydrocodone, fentanyl and methadone [2].

Physicians and other health care providers can make a major contribution toward reducing the toll of opioid overdose through the care they take in prescribing opioid analgesics and monitoring patients' response, as well as their acuity in identifying and effectively addressing opioid overdose. Federally funded CME courses are available at no charge at www.OpioidPrescribing.com (five courses funded by the Substance Abuse and Mental Health Services Administration) and on MedScape (two courses funded by the National Institute on Drug Abuse).

OPIOID OVERDOSE

The risk of opioid overdose can be minimized through adherence to the following clinical practices, which are supported by a considerable body of evidence [3-6].

ASSESS THE PATIENT. Obtaining a history of the patient's past use of drugs (either illicit drugs or prescribed medications with abuse potential) is an essential first step in appropriate prescribing. Such a history should include very specific questions. For example:

- In the past 6 months, have you taken any medications to help you calm down, keep from getting nervous or upset, raise your spirits, make you feel better, and the like?
- Have you been taking any medications to help you sleep? Have you been using alcohol for this purpose?
- Have you ever taken a medication to help you with a drug or alcohol problem?
- Have you ever taken a medication for a nervous stomach?
- Have you taken a medication to give you more energy or to cut down on your appetite?

The patient history also should include questions about use of alcohol and over-the-counter (OTC) preparations. For example, the ingredients in many common cold preparations include alcohol and other central nervous system (CNS) depressants, so these products should not be used in combination with opioid analgesics.

Positive answers to any of these questions warrant further investigation.

TAKE SPECIAL PRECAUTIONS

WITH NEW PATIENTS. Many experts recommend that additional precautions be taken in prescribing for new patients [5,6]. These might involve the following:

1. **Assessment:** In addition to the patient history and examination, the physician should determine who has been caring for the patient in the past, what medications have been prescribed and for what indications, and what substances (including alcohol, illicit drugs and OTC products) the patient has reported using. Medical records should be obtained (with the patient's consent).
2. **Emergencies:** In emergency situations, the physician should prescribe the smallest possible quantity, typically not exceeding a 3 days' supply of an opioid analgesic and arrange for a return visit the next day. At a minimum, the patient's identity should be verified by asking for proper identification.
3. **Limit quantities:** In non-emergency situations, only enough of an opioid analgesic should be prescribed to meet the patient's needs until the next appointment. The patient should be directed to *return to the office* for additional prescriptions, as telephone orders do not allow the physician to reassess the patient's continued need for the medication.

INFORMATION FOR PRESCRIBERS

STATE PRESCRIPTION DRUG MONITORING PROGRAMS (PDMPs)

have emerged as a key strategy for addressing the misuse and abuse of prescription opioids and thus preventing opioid overdoses and deaths. Specifically, prescribers can check their state's PDMP database to determine whether a patient is filling the prescriptions provided and/or obtaining prescriptions for the same or similar drug from multiple physicians.

While many states now have operational PDMPs, the programs differ from state to state in terms of the exact information collected, how soon that information is available to physicians, and who may access the data. Therefore, information about the program in a particular state is best obtained directly from the PDMP or from the state board of medicine or pharmacy.

SELECT AN APPROPRIATE MEDICATION. Rational drug therapy demands that the efficacy and safety of all potentially useful medications be reviewed for their relevance to the patient's disease or disorder [3,6].

When an appropriate medication has been selected, the *dose*, *schedule*, and *formulation* should be determined. These choices often are just as important in optimizing pharmacotherapy as the choice of medication itself. Decisions involve (1) dose (based not only on age and weight of the patient, but also on severity of the disorder, possible loading-dose requirement, and the presence of potentially interacting drugs); (2) timing of administration (such as a bedtime dose to minimize problems associated with sedative or respiratory depressant effects); (3) route of administration (chosen to improve compliance/adherence as well as to attain peak drug concentrations rapidly); and (4) formulation (e.g., selecting a patch in preference to a tablet, or an extended-release product rather than an immediate-release formulation).

Additional safeguards are recommended before prescribing an opioid analgesic. For example, even when sound medical indications have been established, physicians typically consider three additional factors before deciding to prescribe [3,6]:

1. The **severity of symptoms**, in terms of the patient's ability to accommodate them. Relief of symptoms is a legitimate goal of medical practice, but using opioid analgesics requires caution.
2. The patient's **reliability in taking medications**, noted through observation and careful history-taking. The physician should assess a patient's history of and risk factors for drug abuse before prescribing any psychoactive drug and weigh the benefits against the risks. The likely development of physical dependence in patients on long-term opioid therapy should be monitored through periodic check-ups.

3. The **dependence-producing potential of the medication**. The physician should consider whether a product with less potential for abuse, or even a non-drug therapy, would provide equivalent benefits. Patients should be warned about possible adverse effects caused by interactions between opioids and other medications or illicit substances, including alcohol.

At the time a drug is prescribed, patients should be informed that it is illegal to sell, give away, or otherwise share their medication with others, including family members. The patient's obligation extends to keeping the medication in a locked cabinet or otherwise restricting access to it and to safely disposing of any unused supply (visit <http://www.fda.gov/ForConsumers/ConsumerUpdates/ucm101653.htm> for advice from the FDA on how to safely dispose of unused medications).

EDUCATE THE PATIENT AND OBTAIN INFORMED CONSENT.

Obtaining informed consent involves informing the patient about the risks and benefits of the proposed therapy and of the ethical and legal obligations such therapy imposes on both the physician and patient [6]. Such informed consent can serve multiple purposes: (1) it provides the patient with information about the risks and benefits of opioid therapy; (2) it fosters adherence to the treatment plan; (3) it limits the potential for inadvertent drug misuse; and (4) it improves the efficacy of the treatment program.

Patient education and informed consent should specifically address the potential for physical dependence and cognitive impairment as side effects associated with

INFORMATION FOR PRESCRIBERS

opioid analgesics. Other issues that should be addressed in the informed consent or treatment agreement include the following [6]:

- The agreement instructs the patient to stop taking all other pain medications, unless explicitly told to continue by the physician. Such a statement reinforces the need to adhere to a single treatment regimen.
- The patient agrees to obtain the prescribed medication from only one physician and, if possible, from one designated pharmacy.
- The patient agrees to take the medication only as prescribed (for some patients, it may be possible to offer latitude to adjust the dose as symptoms dictate).
- The agreement makes it clear that the patient is responsible for safeguarding the written prescription and the supply of medications, and arranging refills during regular office hours. This responsibility includes planning ahead so as not to run out of medication during weekends or vacation periods.
- The agreement specifies the consequences for failing to adhere to the treatment plan, which may include weaning and discontinuation of opioid therapy if the patient's actions compromise his or her safety.

Both patient and physician should sign the informed consent agreement, and a copy should be placed in the patient's medical record. It also is helpful to give the patient a copy of the agreement to carry with him or her, to document the source and reason for any controlled drugs in his or her possession. Some physicians provide a laminated card that identifies the individual as a patient of their practice. This is helpful to other physicians who may see the patient and in the event the patient is seen in an emergency department.

EXECUTE THE PRESCRIPTION ORDER. Careful execution of the prescription order can prevent manipulation by the patient or others intent on obtaining opioids for non-medical purposes. For example, federal law requires that prescription orders for controlled substances be signed and dated on the day they are issued. Also under federal law, every prescription order must include at least the following information:

- Name and address of the patient
- Name, address and DEA registration number of the physician
- Signature of the physician
- Name and quantity of the drug prescribed
- Directions for use
- Refill information
- Effective date if other than the date on which the prescription was written

Many states impose additional requirements, which the physician can determine by consulting the state medical licensing board. In addition, there are special federal requirements for drugs in different schedules of the federal Controlled Substances Act (CSA), particularly those in Schedule II, where many opioid analgesics are classified.

Blank prescription pads, as well as information such as the names of physicians who recently retired, left the state, or died all can be used to forge prescriptions. Therefore, it is a sound practice to store blank prescriptions in a secure place rather than leaving them in examining rooms.

NOTE: The physician should immediately report the theft or loss of prescription blanks to the nearest field office of the federal Drug Enforcement Administration and to the state board of medicine or pharmacy.

MONITOR THE PATIENT'S RESPONSE TO TREATMENT. Proper prescription practices do not end when the patient receives a prescription. Plans to monitor for drug efficacy and safety, compliance, and potential development of tolerance must be documented and clearly communicated to the patient [3].

Subjective symptoms are important in monitoring, as are objective clinical signs (such as body weight, pulse rate, temperature, blood pressure, and levels of drug metabolites in the bloodstream). These can serve as early signs of therapeutic failure or unacceptable adverse drug reactions that require modification of the treatment plan.

Asking the patient to keep a log of signs and symptoms gives him or her a sense of participation in the treatment

INFORMATION FOR PRESCRIBERS

program and facilitates the physician's review of therapeutic progress and adverse events.

Simply recognizing the potential for non-adherence, especially during prolonged treatment, is a significant step toward improving medication use [7]. Steps such as simplifying the drug regimen and offering patient education also improve adherence, as do phone calls to patients, home visits by nursing personnel, convenient packaging of medication, and periodic urine testing for the prescribed opioid as well as any other respiratory depressant.

Finally, the physician should convey to the patient through attitude and manner that any medication, no matter how helpful, is only part of an overall treatment plan.

When the physician is concerned about the behavior or clinical progress (or the lack thereof) of a patient being treated with an opioid analgesic, it usually is advisable to seek a consultation with an expert in the disorder for which the patient is being treated *and* an expert in addiction. Physicians place themselves at risk if they continue to prescribing opioids in the absence of such consultations [6].

CONSIDER PRESCRIBING NALOXONE ALONG WITH THE PATIENT'S INITIAL OPIOID PRESCRIPTION. With proper education, patients on long-term opioid therapy and others at risk for overdose may benefit from having a naloxone kit to use in the event of overdose [8].

Patients who are candidates for such kits include those who are:

- Taking high doses of opioids for long-term management of chronic malignant or non-malignant pain.
- Receiving rotating opioid medication regimens (and thus at risk for incomplete cross-tolerance).
- Discharged from emergency medical care following opioid intoxication or poisoning.
- At high risk for overdose because of a legitimate medical need for analgesia, coupled with a suspected or confirmed history of substance abuse, dependence, or non-medical use of prescription or illicit opioids.
- Completing mandatory opioid detoxification or abstinence programs.
- Recently released from incarceration and a past user or abuser of opioids (and presumably with reduced opioid tolerance and high risk of relapse to opioid use).

It also may be advisable to suggest that the at-risk patient create an "overdose plan" to share with friends, partners and/or caregivers. Such a plan would contain information on the signs of overdose and how to administer naloxone or otherwise provide emergency care (as by calling 911).

DECIDE WHETHER AND WHEN TO END OPIOID THERAPY.

Certain situations may warrant immediate cessation of prescribing. These generally occur when out-of-control behaviors indicate that continued prescribing is unsafe or causing harm to the patient [3]. Examples include altering or selling prescriptions, accidental or intentional overdose, multiple episodes of running out early (due to excessive use), doctor shopping, or threatening behavior.

When such events arise, it is important to separate the patient as a person from the behaviors caused by the disease of addiction, as by demonstrating a positive regard for the person but no tolerance for the aberrant behaviors.

The essential steps are to (1) stop prescribing, (2) tell the patient that continued prescribing is not clinically supportable (and thus not possible), (3) urge the patient to accept a referral for assessment by an addiction specialist, (4) educate the patient about signs and symptoms of spontaneous withdrawal and urge the patient to go to the emergency department if symptoms occur, and (5) assure the patient that he or she will continue to receive care for the presenting symptoms or condition [6].

INFORMATION FOR PRESCRIBERS

Identification of a patient who is abusing a prescribed controlled drug presents a major therapeutic opportunity. The physician should have a plan for managing such a patient, typically involving work with the patient and the patient's family, referral to an addiction expert for assessment and placement in a formal addiction treatment program, long-term participation in a 12-Step mutual help program such as Narcotics Anonymous, and follow-up of any associated medical or psychiatric comorbidities [3].

In all cases, patients should be given the benefit of the physician's concern and attention. It is important to remember that even drug-seeking patients often have very real medical problems that demand and deserve the same high-quality medical care offered to any patient [3,6].

TREATING OPIOID OVERDOSE

In the time it takes for an overdose to become fatal, it is possible to reverse the respiratory depression and other effects of opioids through respiratory support and administration of the opioid antagonist naloxone (Narcan) [9]. Naloxone is approved by the FDA and has been used for decades to reverse overdose and resuscitate individuals who have overdosed on opioids.

The safety profile of naloxone is remarkably high, especially when used in low doses and titrated to effect [8,9]. If given to individuals who are not opioid-intoxicated or opioid-dependent, naloxone produces no clinical effects, even at high doses. Moreover, while rapid opioid withdrawal in tolerant patients may be unpleasant, it is not typically life-threatening.

Naloxone should be part of an overall approach to opioid overdose that incorporates the following steps.

RECOGNIZE THE SIGNS OF OVERDOSE. An opioid overdose requires rapid diagnosis. The most common signs of overdose include [3]:

- Pale and clammy face
- Limp body
- Fingernails or lips turning blue/purple
- Vomiting or gurgling noises
- Cannot be awakened from sleep or is unable to speak
- Very little or no breathing
- Very slow or no heartbeat

Signs of **OVERMEDICATION**, which may progress to overdose, include [3]:

- Unusual sleepiness or drowsiness
- Mental confusion, slurred speech, intoxicated behavior
- Slow or shallow breathing
- Pinpoint pupils
- Slow heartbeat, low blood pressure
- Difficulty waking the individual from sleep

Because opioids depress respiratory function and breathing, one telltale sign of an individual in a critical medical state is the "death rattle." Often mistaken for snoring, the "death rattle" is an exhaled breath with a very distinct, labored sound coming from the throat. It indicates that emergency resuscitation is needed immediately [8].

SUPPORT RESPIRATION. Supporting respiration is the single most important intervention for opioid overdose and may be life-saving on its own. Ideally, individuals who are experiencing opioid overdose should be ventilated with 100% oxygen before naloxone is administered to reduce the risk of acute lung injury [3, 8]. In situations where 100% oxygen is not available, rescue breathing can be very effective in supporting respiration [8]. Rescue breathing involves the following steps:

- Verify that the airway is clear.
- With one hand on the patient's chin, tilt the head back and pinch the nose closed.
- Place your mouth over the patient's mouth to make a seal and give 2 slow breaths (the patient's chest should rise, but not the stomach).
- Follow up with one breath every 5 seconds.

INFORMATION FOR PRESCRIBERS

ADMINISTER NALOXONE. Naloxone (Narcan) should be given to any patient who presents with signs of opioid overdose, or when overdose is suspected [8]. Naloxone can be given by intramuscular or intravenous injection every 2 to 3 minutes [8-10].

The most rapid onset of action is achieved by intravenous administration, which is recommended in emergency situations [9]. Intravenous administration generally is used with patients who have no history of opioid dependence. Opioid-naive patients may be given starting doses of up to 2 mg without concern for triggering withdrawal symptoms [8].

The intramuscular route of administration may be more suitable for patients with a history of opioid dependence because it provides a slower onset of action and a prolonged duration of effect, which may minimize rapid onset of withdrawal symptoms [8].

Pregnant patients. Naloxone can safely be used to manage opioid overdose in pregnant women. The lowest dose to maintain spontaneous respiratory drive should be used to avoid triggering acute opioid withdrawal, which may cause fetal distress [8].

MONITOR THE PATIENT'S RESPONSE. Patients should be monitored for re-emergence of signs and symptoms of opioid toxicity for at least 4 hours following the last dose of naloxone (however, patients who have overdosed on long-acting opioids require more prolonged monitoring) [8].

Most patients respond to naloxone by returning to spontaneous breathing, with mild withdrawal symptoms [8]. The response generally occurs within 3 to 5 minutes of naloxone administration. (Rescue breathing should continue while waiting for the naloxone to take effect.)

The duration of effect of naloxone is 30 to 90 minutes. Patients should be observed after that time for re-emergence of overdose symptoms. The goal of naloxone therapy should be restoration of adequate spontaneous breathing, but not necessarily complete arousal [8-10].

More than one dose of naloxone may be required to revive the patient. Those who have taken longer-acting opioids may require further intravenous bolus doses or an infusion of naloxone [8]. Therefore, it is essential to get the person to an emergency department or other source of acute care as quickly as possible, even if he or she revives after the initial dose of naloxone and seems to feel better.

SIGNS OF OPIOID WITHDRAWAL: Withdrawal triggered by naloxone can feel unpleasant. As a result, some persons become agitated or combative when this happens and need help to remain calm.

The signs and symptoms of opioid withdrawal in an individual who is physically dependent on opioids may include, but are not limited to, the following: body aches, diarrhea, tachycardia, fever, runny nose, sneezing, piloerection, sweating, yawning, nausea or vomiting, nervousness, restlessness or irritability, shivering or trembling, abdominal cramps, weakness, and increased blood pressure [9]. Withdrawal syndromes may be precipitated by as little as 0.05 to 0.2 mg intravenous naloxone in a patient taking 24 mg per day of methadone.

In neonates, opioid withdrawal also may produce convulsions, excessive crying, and hyperactive reflexes [9].

NALOXONE-RESISTANT PATIENTS: If a patient does not respond to naloxone, an alternative explanation for the clinical symptoms should be considered. The most likely explanation is that the person is not overdosing on an opioid but rather some other substance or may even be experiencing a non-overdose medical emergency. A possible explanation to consider is whether the individual has overdosed on buprenorphine, a long-acting opioid partial agonist. Because buprenorphine has a higher affinity for the opioid receptors than do other opioids, naloxone may not be effective at reversing the effects of buprenorphine-induced opioid overdose [8].

In all cases, support of ventilation, oxygenation, and blood pressure should be sufficient to prevent the complications of opioid overdose and should be given the highest priority if the patient's response to naloxone is not prompt.

NOTE: All naloxone products have an expiration date. It is important to check the expiration date and obtain replacement naloxone as needed.

INFORMATION FOR PRESCRIBERS

LEGAL AND LIABILITY CONSIDERATIONS

Health care professionals who are concerned about legal risks associated with prescribing naloxone may be reassured by the fact that prescribing naloxone to manage opioid overdose is consistent with the drug's FDA-approved indication, resulting in no increased liability so long as the prescriber adheres to general rules of professional conduct. State laws and regulations generally prohibit physicians from prescribing a drug such as naloxone to a third party, such as a caregiver. (Illinois, Massachusetts, New York, and Washington State are the exceptions to this general principle.) More information on state policies is available at www.prescribetoprevent.org/ or from individual state medical boards.

CLAIMS CODING AND BILLING

Most private health insurance plans, Medicare, and Medicaid cover naloxone for the treatment of opioid overdose, but policies vary by state. The cost of take-home naloxone should not be a prohibitive factor. Not all community pharmacies stock naloxone routinely but can always order it. If you are caring for a large patient population likely to benefit from naloxone you may wish to notify the pharmacy when you implement naloxone prescribing as a routine practice.

The codes for Screening, Brief Intervention, and Referral to Treatment (SBIRT) can be used to bill time for counseling a patient about how to recognize overdose and how to administer naloxone. Billing codes for SBIRT are as follows:

Commercial Insurance: CPT 99408 (15 to 30 minutes)

Medicare: G0396 (15 to 30 minutes)

Medicaid: H0050 (per 15 minutes)

RESOURCES FOR PRESCRIBERS

Additional information on prescribing opioids for chronic pain is available at the following websites:

www.opioidprescribing.com. Sponsored by the Boston University School of Medicine, with support from SAMHSA, this site presents course modules on various aspects of prescribing opioids for chronic pain. To view the list of courses and to register, go to <http://www.opioidprescribing.com/overview>. CME credits are available at no charge.

www.pcss-o.org or www.pcssb.org.

Sponsored by the American Academy of Addiction Psychiatry in collaboration with other specialty societies and with support from SAMHSA, the Prescriber's Clinical Support System offers multiple resources related to opioid prescribing and the diagnosis and management of opioid use disorders.

www.medscape.com. Two course modules sponsored by the National Institute on Drug Abuse and posted on MedScape can be accessed at <http://www.medscape.org/viewarticle/770687> and <http://www.medscape.org/viewarticle/770440>. CME credits are available.

REFERENCES

1. Centers for Disease Control and Prevention (CDC). CDC grand rounds: Prescription drug overdoses — A U.S. epidemic. *MMWR Morb Mortal Wkly Rep.* 2012;61(1):10–13.
2. Harvard Medical School. Painkillers fuel growth in drug addiction; Opioid overdoses now kill more people than cocaine or heroin. *Harvard Ment Hlth Let.* 2011;27(7):4–5.
3. Beletsky L, Rich JD, Walley AY. Prevention of fatal opioid overdose. *JAMA.* 2012 Nov 14;308(18):1863–1864.
4. Isaacson JH, Hopper JA, Alford DP, Parran T. Prescription drug use and abuse. Risk factors, red flags, and prevention strategies. *Postgrad Med.* 2005;118:19.
5. Coffin PO, Sullivan SD. Cost-effectiveness of distributing naloxone to heroin users for lay overdose reversal. *Ann Internal Med.* 2013;58:1–9.
6. Finch JW, Parran TV, Wilford BB, Wyatt SA. Clinical, legal and ethical considerations in prescribing drugs with abuse potential. In Ries RK, Alford DP, Saitz R, Miller S, eds. *Principles of Addiction Medicine, Fifth Edition.* Philadelphia, PA: Lippincott, Williams & Wilkins, Ch. 109, in press 2013.
7. Michna E, Ross EL, Hynes WL, et al. Predicting aberrant drug behavior in patients treated for chronic pain: Importance of abuse history. *J Pain Symptom Manage.* 2004;28:250.
8. BMJ Evidence Centre. Treatment of opioid overdose with naloxone. *British Medical Journal.* Updated October 23, 2012. [Accessed March 24, 2013, at www.bmj.com]
9. Rx List [Accessed March 24, 2013, at www.rxlist.com]
10. Drugs.com [Accessed March 24, 2013, at www.drugs.com]

Acknowledgments

This publication was prepared for the Substance Abuse and Mental Health Services Administration (SAMHSA) by the Association of State and Territorial Health Officials, in cooperation with Public Health Research Solutions, under contract number 10-233-00100 with SAMHSA, U.S. Department of Health and Human Services (HHS). LCDR Brandon Johnson, M.B.A., served as the Government Project Officer.

Disclaimer

The views, opinions, and content of this publication are those of the authors and do not necessarily reflect the views, opinions, or policies of SAMHSA or HHS.

Public Domain Notice

All materials appearing in this volume except those taken directly from copyrighted sources are in the public domain and may be reproduced or copied without permission from SAMHSA or the authors. Citation of the source is appreciated. However, this publication may not be reproduced or distributed for a fee without the specific, written authorization of the Office of Communications, SAMHSA, HHS.

Electronic Access and Copies of Publication

This publication may be ordered from SAMHSA's Publications Ordering Web page at www.store.samhsa.gov. Or, please call SAMHSA at 1-877-SAMHSA-7 (1-877-726-4727) (English and Español).

Recommended Citation

Substance Abuse and Mental Health Services Administration. SAMHSA Opioid Overdose Prevention Toolkit: Information for Prescribers. HHS Publication No. (SMA) 13-4742. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2013.

Originating Office

Division of Pharmacologic Therapies, Center for Substance Abuse Treatment, Substance Abuse and Mental Health Services Administration, 1 Choke Cherry Road, Rockville, MD 20857.



HHS Publication No. (SMA) 13-4742
Printed 2013